



MYCORRHIZA BIOFERTILIZER FOR SUSTAINABLE AGRICULTURE

■ K.R.K Prasad, Monika Singh and Ashwani Kumar

¹Division of Crop Improvement, Central Soil Salinity Research Institute, Karnal (HARYANA) INDIA
Email : ankita.garcoti2010@gmail.com

An arbuscular mycorrhiza is a type of mycorrhiza in which the fungus penetrates the cortical cells of the roots of a vascular plant. Arbuscular mycorrhizae (AMs) are characterized by the formation of unique structures such as arbuscules and vesicles by fungi of the phylum Glomeromycota (AM fungi). AM fungi (AMF) help plants to capture nutrients such as phosphorus and micronutrients from the soil. It is believed that the development of the arbuscular mycorrhizal symbiosis played a crucial role in the initial colonisation of land by plants and in the evolution of the vascular plants. Spores of the AM fungi are thick walled multi-nucleate resting structures. There are three major groups of mycorrhiza: Ectomycorrhiza, Ectendomycorrhiza and Endomycorrhiza. Ectomycorrhiza and endomycorrhiza are important in agriculture and forestry. Endomycorrhiza (vesicular arbuscular mycorrhiza; VA mycorrhiza; now known as arbuscular mycorrhiza, AM) play a very important role on enhancing the plant growth and yield due to an increase supply of phosphorus to the host plant. Mycorrhizal plants can absorb and accumulate several times more phosphate from the soil or solution than non-mycorrhizal plants. Plants inoculated with endomycorrhiza have been shown to be more resistant to some root diseases. It is now generally recognized that they improve not only the phosphorus nutrition of the host plant but also its growth, which may result in an increase in resistance to drought stress and some diseases. Therefore, AM fungi offer a great potential for sustainable agriculture, and the application of AM fungi to agriculture has been developed.

Nutrient uptake and exchange:

An increase in the carbon supplied by the plant to the AM fungi increases the uptake of phosphorus and the transfer of phosphorus from fungi to plant (Bücking & Shachar-Hill

2005). Phosphorus uptake and transfer is also lowered when the photosynthate supplied to the fungi is decreased. Species of AMF differ in their abilities to supply the plant with phosphorus. In some cases arbuscular mycorrhizae are poor symbionts providing little phosphorus while taking relatively high amounts of carbon

The benefit of mycorrhizae to plants is mainly attributed to increased uptake of nutrients, especially phosphorus. This increase in uptake may be due to increase surface area of soil contact, increased movement of nutrients into mycorrhizae, a modification of the root environment and increased storage. Mycorrhizae can be much more efficient than plant roots at taking up phosphorus. Phosphorus travels to the root or via diffusion and hyphae reduce the distance required for diffusion thus increasing uptake. The rate of inflow of phosphorus into mycorrhizae can be up to six times that of the root hairs. In some cases the role of phosphorus uptake can be completely taken over by the mycorrhizal network and all of the plant's phosphorus may be of hyphal origin.

Benefits of Mycorrhizal Biofertilizer :

Mycorrhiza plays a very important role on enhancing the plant growth and yield due to an increase supply of phosphorus to the host plant. Mycorrhizal plants can absorb and accumulate several times more phosphate from the soil or solution than non-mycorrhizal plants. Plants inoculated with endomycorrhiza have been shown to be more resistant to some root diseases.

Mycorrhiza increase root surface area for water and nutrients uptake. The use of mycorrhizal biofertilizer helps to improve higher branching of plant roots, and the mycorrhizal hyphae grow from the root to soil enabling the plant roots to contact with wider area of soil surface, hence, increasing the absorbing area for water and nutrients absorption of the plant

root system. Therefore, plants with mycorrhizal association will have higher efficiency for nutrients absorption, such as nitrogen, phosphorus, potassium, calcium, magnesium, zinc, and copper; and also increase plant resistance to drought.

Benefits of mycorrhizal biofertilizer as follows:

- Allow plants to take up nutrients in unavailable forms or nutrients that are fixed to the soil. Some plant nutrients, especially phosphorus, are elements that dissolve were in water in neutral soil. In the extreme acidic or basic soil, phosphorus is usually bound to iron, aluminum, calcium, or magnesium, leading to water insolubility, which is not useful for plants.
- Mycorrhiza plays an important role in phosphorus absorption for plant via cell wall of mycorrhiza to the cell wall of plant root. In addition, mycorrhiza help to absorb other organic substances that are not fully soluble for plants to use, and also help to absorb and dissolve other nutrients for plants by storage in the root.
- Enhance plant growth, improve crop yield, and increase income for the farmers. Arising from improved water and essential nutrients absorption for plant growth by mycorrhiza, it leads to improvement in plant photosynthesis, nutrients translocation, and plant metabolism processes.
- As in the trial involving mycorrhizal biofertilizer on asparagus it was observed that, when the farmers used suggested amount of chemical fertilizer together with mycorrhizal biofertilizer, it was found that the crop yield improved by more than 50%, and the farmers' income increased 61% higher than when chemical fertilizer alone was used.
- Improve plant resistance to root rot and collar rot diseases. Mycorrhizal association in plant roots will help plant to resist root rot and collar rot diseases caused by other fungi.
- It can be used together with other agricultural chemicals. Mycorrhiza are endurable to several chemical substances; for example; pesticide such as endrin, chlordane, methyl parathion, methomyl

carbofuran; herbicide such as glyphosate, fuazifopbutyl; chemical agents for plant disease elimination such as captan, benomyl, maneb triforine and mancozeb .

Therefore, the plant has better growth and yield, reduce the use of chemical fertilizer, sometimes up to half of the suggested amount, which in turn increases income for the farmers.

In fact, in some countries the AM fungal inocula have been commercialized. Since it is laborious and cost-consuming for production of AM fungal inocula because of their obligate bio tropic nature, the ways to increase the function of the indigenous AM fungi in soil have also been developed.

REFERENCES

- Ames R N, Reid C P P, Porter L K and Cambardella C (1983)** Hyphal uptake and transport of nitrogen from two ¹⁵N labelled sources by *Glomus mosseae*, a vesicular-arbuscular mycorrhizal fungus. *New Phytol.* **95**, 381–396
- Bolan NS (1991).** A critical review of the role of mycorrhizal fungi in the uptake of phosphorus by plants. *Plant Soil* **134**, 189–207
- Brundrett, M. et al.** Working with Mycorrhizas in Forestry and Agriculture, Australian Centre for International Agricultural Research (1996).
- International Culture Collection of Arbuscular and Vesicular-Arbuscular Mycorrhizal Fungi <http://invam.caf.wvu.edu>
- Saito M and Marumoto T (2002).** Inoculation with arbuscular mycorrhizal fungi: the status quo in Japan and the future prospects. *Plant Soil* **244**: 273-279
- Trappe J M (1987).** “Phylogenetic and ecologic aspects of mycotrophy in the angiosperms from an evolutionary standpoint”. *Ecophysiology of VA Mycorrhizal Plants*, G.R. Safir (EDS)(Florida: CRC Press).
- Walker C and Vestberg M (1994).** A simple and inexpensive method for producing and maintaining closed pot cultures of arbuscular mycorrhizal fungi. *Agricultural Science in Finland* **3**: 233-240.

Conventional agriculture plays a significant role in meeting the food demands of a growing human population, which has also led to an increasing dependence on chemical Fertilizers and pesticides [1]. Chemical fertilizers are industrially manipulated, substances composed of known quantities of nitrogen, phosphorus and potassium, and their exploitation causes air and ground water pollution by eutrophication of water bodies [2]. In this regard

• Mechanism of action of various biofertilizers. Mycorrhiza is the association of fungus with the roots of higher plants.

• Raja N: Biopesticides and biofertilizers: ecofriendly sources for sustainable agriculture. *J Biofertil Biopestici*. 2013, 1000e112: 1000e112 potential tool for sustainable agriculture. The mycorrhizal symbiosis becomes even more important in sustainable agricultural systems where nutrient inputs are low.

• Their potential as a biofertilizer lies in their mycorrhizal benefits and plant-soil interactions. Hence, their selection for inoculum and

• Arbuscular Mycorrhizas on Sustainable Agriculture and Natural Ecosystems. BirkhÓuser Verlag, Basel, Switzerland, pp. 201-215. The application of arbuscular mycorrhizal fungi towards ensuring sustainable agriculture is gaining more awareness each day with their role in improving plant health, productivity, and soil fertility.

5.5. Sulphur Oxidizing Microbes. Sulphur is an essential macronutrient needed by plants in high concentration.

• Yadav, K.K.; Sarkar, S. Biofertilizers, Impact on Soil Fertility and Crop Productivity under Sustainable Agriculture. *Environ. Ecol. Biofertilizers for Sustainable Agriculture and Environment* pp 501-520 | Cite as. The Role of Arbuscular Mycorrhiza in Sustainable Environment and Agriculture. Authors. Authors and affiliations.

• Arbuscular mycorrhizal (AM) fungi are a class of beneficial microorganisms that are widely distributed in soil ecosystems and can form symbiotic associations with more than 90% of terrestrial higher plants. They play an important role in promoting plant growth, improving plant disease resistance and stress resistance, and maintaining the sustainable development of agricultural ecosystem.