

- folate, and homocysteine to cognitive performance in the Normative Aging Study. *Am J Clin Nutr* 1996;63:306
18. McCaddon A, Davies G, Hudson P, et al. Total serum homocysteine in senile dementia of the Alzheimer's type. *Int J Geriatr Psychiatry* 1998;13:235
 19. Blijhan GH. Homocysteine metabolism 2nd international conference. *Neth J Med* 1998;52(suppl):S1
 20. Miller JW, Green R, Allen LH, Mungas DM, Haan MN. Homocysteine correlates with cognitive function in the Sacramento Area Latino Study of Aging (SALSA). *FASEB J* 1999;13:A374
 21. Folstein MF, Folstein FE, McHugh PR. Mini-mental state: a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189
 22. Kalmijn S, Launer LJ, Lindemans J, et al. Total homocysteine and cognitive decline in a community-based sample of elderly subjects: the Rotterdam Study. *Am J Epidemiol* 1999;150:283
 23. Selhub J, Jacques PF, Wilson PWF, Rush D, Rosenberg IH. Vitamin status and intake as primary determinants of homocysteinemia in an elderly population. *JAMA* 1993;270:2693
 24. Jacques PF, Selhub J, Bostom AG, Wilson PW, Rosenberg IH. The effect of folic acid fortification on plasma folate and total homocysteine concentrations. *N Engl J Med* 1999;340:1449
 25. Lawrence JM, Petitti DB, Watkins M, Umekubo MA. Trends in serum folate after food fortification. *Lancet* 1999;354:915
 26. Green R, Miller JW. Folate deficiency beyond megaloblastic anemia: hyperhomocysteinemia and other manifestations of dysfunctional folate metabolism. *Semin Hematol* 1999;36:47
 27. Chanarin I. *The Megaloblastic Anemias, 2nd ed.* Oxford: Blackwell Scientific Publications, 1979
 28. Carmel R. Prevalence of undiagnosed pernicious anemia in the elderly. *Arch Intern Med* 1996;156:1097
 29. Carmel R, Sinow RM, Siegel ME, et al. Food cobalamin malabsorption occurs frequently in patients with unexplained low serum cobalamin levels. *Arch Int Med* 1988;148:1715
 30. Pennypacker L, Allen RH, Kelly JP, et al. High prevalence of cobalamin deficiency in elderly outpatients. *J Am Geriatr Soc* 1991;39:1155
 31. Kuzminski AM, Del Giacco EJ, Allen RH, Stabler SP, Lindenbaum J. Effective treatment of cobalamin deficiency with oral cobalamin. *Blood* 1998;92:1191
 32. Miller JW, Green R, Herbert VD, Flynn MA. Oral vitamin B₁₂ supplementation decreases homocysteine in healthy elderly people with suboptimal vitamin B₁₂ status. *FASEB J* 1999;13:A936

Chicory Fructooligosaccharides and the Gastrointestinal Tract

Marcel B. Roberfroid, PhD

From the Université Catholique de Louvain, Brussels, Belgium

Chicory fructooligosaccharides are fructans extracted on a commercial basis from the chicory root. These materials are called *inulin* or *oligofructose*, and they are widely used in Europe in a variety of foods, especially as dietary fiber. This editorial examines the composition and source of inulin and oligofructose, the physiologic effects of their consumption, and how these materials relate to the concept of dietary fiber.

CHICORY FRUCTOOLIGOSACCHARIDES: IDENTITY AND OCCURRENCE

Chicory fructooligosaccharides are fructans. *Fructan* is a general term used for any carbohydrate in which fructosyl-fructose links constitute the majority of glycosidic bonds. The plant species currently used by the food industry to produce inulin belongs to the Compositae family; it is chicory (*Cichorium intybus*).¹ In chicory inulin, both GpyFn (α -D-glucopyranosyl-[β -D-fructofuranosyl]*n*-1-D-fructofuranoside) and FpyFn (β -D-fructopyranosyl-[β -D-fructofuranosyl]*n*-1-D-fructofuranoside) compounds are considered to be included under the same nomenclature, and the number of fructose units differs from 2 to more than 70. Native inulin is processed by the food industry to produce either short-chain fructans, namely oligofructose (DP 2–10; average DP 4), as a result of partial enzymatic (endo-inulinase EC 3.2.1.7) hydrolysis or long-chain fructans by applying industrial physical-separation techniques. Chicory fructooligosaccharides are present in significant amounts in several edible fruits and vegetables. Average daily consumption has been estimated to be between 1 and 4 g in the United States³ and between 3 and 11 g in Europe,² the most common sources being wheat, onion, banana, garlic, and leek.²

Chicory inulin and oligofructose are officially recognized as natural food ingredients and are classified as dietary fiber in almost all European countries. An analytical method has recently been developed to quantify inulin and oligofructose in plants and food products.⁴

METABOLISM AND PHYSIOLOGIC EFFECTS OF CHICORY FRUCTOOLIGOSACCHARIDES

Non-Digestibility in the Upper Gastrointestinal Tract

Because of the β -configuration of the anomeric C2 in their fructose monomers, inulin and oligofructose are resistant to hydrolysis by human digestive enzymes (α -glucosidase, maltase-isomaltase, sucrase) specific for α -glycosidic linkages. These carbohydrates have indeed been classified as "non-digestible" oligosaccharides.^{5,6} Both in vitro^{7,8} and in vivo^{9–11} data have supported this classification. Knudsen and Hesso¹⁰ and Ellegård et al.¹¹ used the ileostomy model, and both studies have shown that between 86% and 88% of the ingested dose (10, 17, or 30 g) of inulin and oligofructose are recovered in the ileostomy effluent, supporting the conclusion that chicory fructooligosaccharides are practically indigestible in the small intestine of humans. Moreover, there is no evidence that they are absorbed to any significant extent.^{9,12,13} Thus, it has been proposed to classify them as "colonic food," i.e., a "food entering the colon and serving as substrate for the endogenous bacteria, thus indirectly providing the host with energy, metabolic substrates and essential micronutrients."¹⁴

Fermentation in the Large Bowel: The Prebiotic Effect

That inulin and oligofructose are fermented by bacteria colonizing the large bowel is supported by a large number of in vitro and in vivo studies that also have confirmed the production of lactic and short-chain carboxylic acids as endproducts of the fermenta-

Correspondence to: Marcel B. Roberfroid, PhD, Emeritus Professor, Université Catholique de Louvain, 71 rue du COQ, B-1180 Brussels, Belgium. E-mail:roberfroid@pmnt.ucl.ac.be

tion.^{15,16} Furthermore, human *in vivo* studies have demonstrated that this fermentation leads to the selective stimulation of growth of the bifidobacteria population, making inulin and oligofructose the prototype prebiotic.^{15,17–19} Compared with most other malabsorbed carbohydrates (e.g., resistant starch and other dietary fibers), the colonic fermentation of chicory fructooligosaccharides is accompanied by a significant change in the composition of the colonic microbiota due to selective proliferation of bifidobacteria and a concomitant reduction in the number of other bacteria such as bacteroides, fusobacteria, or clostridia. This effect has been identified as “a prebiotic effect” and a prebiotic has been defined as “a non-digestible food ingredient that beneficially affects the host by selectively stimulating the growth and/or the activity of one or a limited number of bacteria in the colon and thus improves host health.”¹⁴ Based on the results of well-designed human studies that have shown significant changes in the composition of human fecal flora, it can be concluded that chicory fructooligosaccharides (3 to 15 g/d for a few weeks) are prebiotic.^{20–22} However, even though some studies have shown a significant reduction in the number of clostridia, the health benefits (e.g., reducing the risk of intestinal infections) of such a change in the composition of the colonic microbiota still need to be established. A recent report by Hunter et al. has shown that, at a daily dose of 6 g (2×3 g), oligofructose had no therapeutic value in patients with irritable bowel syndrome.²³ Conversely, in an experimental model of necrotizing enterocolitis in quails, Catala et al. reported data that support the hypothesis that oligofructose may prevent the overgrowth of bacteria known to play a role in this pathology in preterm neonates.²⁴

The colonic fermentation of inulin and oligofructose produces short-chain fatty acids and lactate plus gases as products of the anaerobic metabolism. Concerning the pattern of production of short-chain fatty acids, *in vitro* fermentation and animal studies have demonstrated that supplementing the diet with inulin and oligofructose decreases the cecal pH and increases the size of the cecal pool of short-chain fatty acids. Due to the selective fermentation of these chicory fructans, the relative proportion of the three main short-chain fatty acids is changed. Typically, the molar ratio of butyrate is increased as compared with rats that received a typical starch-rich standard diet without added oligosaccharides.^{25–29} Possibly related to this increase in the pool of short-chain fatty acids is the effect of inulin and oligofructose on the intestinal tissue, leading to hyperplasia of the mucosa and increased wall thickness in both the small intestine²⁹ and the cecum.²⁷

Other Physiologic Effects

Besides the properties reviewed above concerning their role as fiber in the diet, inulin and oligofructose have been shown to induce interesting physiologic and nutritional effects. These effects relate to improved calcium bioavailability, the reduction of risk in developing precancerous lesions in the colon, hypotriacylglycerolemia, and hypoinsulinemia in experimental models. These effects have been reviewed recently.^{30,31}

One of the most promising physiologic consequences of the consumption of chicory fructooligosaccharides is an increased bioavailability of calcium. Such an effect has extensively been studied in rats and hamsters.^{32–34} These studies have led to the conclusion that, most probably because of their malabsorption and their colonic fermentation, these food ingredients facilitate calcium absorption from the large bowel compartment, thus complementing the process that takes place in the small bowel. Change in colonic pH, production of short-chain fatty acids, and increase in mucosal concentration of the calbindin protein in the colon have been proposed as hypotheses to explain that effect. In addition to increased calcium bioavailability, it has been shown in both rat and hamster that feeding *fructooligosaccharides* increases calcium concentration and improves structure in the bones.^{33,34} More recently,

two human trials (one in *adolescents* and one in *adults*) have shown that supplementing the diet with either 16.8 g of oligofructose³⁵ or 40 g of inulin³⁶ significantly increases the apparent absorption of calcium by 12% and 11%, respectively. The first study used the calcium-balance measurement, whereas the second used the double-stable isotope technique.

Over the past 2 y, reports have been published that have repeatedly demonstrated that feeding chicory fructooligosaccharides to rats previously treated with a colon carcinogen (i.e., dimethylhydrazine or azoxymethane) reduces the incidence of the so-called aberrant crypt foci in the colon.^{37,38} In one of these studies, the synbiotic approach that combines oligofructose (prebiotic) and bifidobacteria (probiotic) was reported to be more active than either the prebiotic or the probiotic alone.³⁸ Even though still experimental, these data suggest that chicory fructooligosaccharides may play a role in reducing the risk of developing preneoplastic lesions in the colon.

With regard to the effect of chicory fructooligosaccharides on the metabolism of triacylglycerols, experimental data have led to formulation of the hypothesis that feeding such non-digestible oligosaccharides may reduce the hepatic-lipogenic capacity by inhibiting the expression of lipogenic enzymes either by the production of short-chain fatty acids or by modulating insulinemia through still-unknown mechanisms that are presently under extensive investigation.³⁰

CALORIC VALUE

Several studies have estimated the caloric value of inulin and oligofructose in human subjects using various methodologies. In summary, because only a part of the energy of these dietary carbohydrates is salvaged, their available energy content is only 40–50% that of a digestible carbohydrate, for an energy value of 1.5 to 2.0 kcal/g.³⁹

SAFETY

Inulin and oligofructose are natural food ingredients present in edible plants and are part of the traditional diet. Based on data from the U.S. Department of Agriculture, it was estimated that the average per-capita daily intake of chicory fructooligosaccharides ranges between 1 and 4 g/d, with the 90th percentile consuming 2 to 8 g/d.³ Their current and proposed uses in multiple food categories could lead, in principle, to increased consumer exposure to dietary fiber in general and to inulin and oligofructose in particular. This may have some desirable effects on colonic function and health. The daily reference value for fiber (soluble or insoluble) is 25 g/d. The average daily intake of inulin or oligofructose, naturally present in food, is estimated to be 1 to 4 g/d in the United States³ and 3 to 11 g/d in Europe.² The extensive available data indicate that safety, as traditionally defined, is not an issue in the case of inulin and oligofructose.⁴⁰ This notion is supported by the critical review of the toxicologic studies showing that chicory fructooligosaccharides do not increase morbidity or mortality or cause reproductive or target-organ toxicity. These fibers are not mutagenic, carcinogenic, or teratogenic.^{40,41}

Numerous animal and human investigational studies have been performed to identify the physiologic and potential health benefits of these fructans and to assess possible intolerance. The only biological effects observed have been attributed to their action as non-digestible, fermentable carbohydrates causing self-limited gastrointestinal distress. Symptoms range from flatulence (the most common), to borborygmi and bloating, to laxation (the most severe). As demonstrated by various studies, the ensuing gastrointestinal symptoms are dose dependent.^{42,43} Results indicate that these fructans are well tolerated at amounts up to 20 g/d; diarrhea can develop with intakes of 30 g/d or more.⁴⁴

CONCLUSION: CHICORY FRUCTOOLIGOSACCHARIDES AS DIETARY FIBER

As is the case with other dietary fibers, inulin and oligofructose are resistant to digestion in the upper part of the intestinal tract and are subsequently fermented in the colon. They also have a bulking effect due to the increase in the microbial biomass that results from their fermentation. From a quantitative point of view, that bulking effect, expressed as the increase in daily fecal mass, has been reported to range between 1.5 and 2 g/d of ingested inulin or oligofructose,^{17,31,44} which is of the same order of magnitude as pectin. Another typical dietary fiber effect is the increase in stool frequency, as was recently observed when inulin was incorporated into the diet of healthy but chronically, slightly constipated human volunteers.^{19,44} Thus, inulin and oligofructose fit well within the current concept of those classes of materials referred to as dietary fiber. Their nutritional properties agree very well with the accepted definition of dietary fiber put forth by Trowell et al.⁴⁵ None of the molecules of fructose and glucose that form inulin and oligofructose appears in the portal blood. These materials share the basic common characteristics of dietary fibers, i.e., saccharides of plant origin, resistance to digestion, absorption in the small intestine, and fermentation in the colon to produce short-chain fatty acids that are absorbed and metabolized in different parts of the body. Moreover, this fermentation induces a bulking effect. Chicory fructooligosaccharides also show beneficial effects on calcium absorption, on the biochemical mechanisms controlling triacylglycerol metabolism, and possibly on the reduction of risk in developing precancerous lesions in the colon.

REFERENCES

- De Bruyn A, Alvarez AP, Sandra P, DeLeenheer L. Isolation and identification of *O*- β -D-fructofuranosyl-(2 \rightarrow 1)-*O*- β -D-fructosuranosyl-(2 \rightarrow 1)-D-fructose: a product of the enzymic hydrolysis of the inulin from *Cichorium intybus*. *Carbohydrate Res* 1992;235:303
- Van Loo J, Coussemont P, De Leenheer L, Hoebregs H, Smits G. On the presence of inulin and oligofructose as natural ingredients in the Western diet. *Crit Rev Food Sci Nutr* 1995;35:525
- MoshFegh AJ, Friday JE, Goldman JP, Chug Ahuga JK. Presence of inulin, and oligofructose in the diets of Americans. *J Nutr* 1999;129(suppl 7S):1407S
- Quemener B, Thibault J, Coussment P. Determination of inulin and oligofructose in food products, and integration in the AOAC method for measurement of total dietary fibre. *Lebensm Wiss U Technol* 1994;27:125
- Cummings JH, Roberfroid MB, Andersson H, et al. A new look at dietary carbohydrate: chemistry, physiology and health. *Eur J Clin Nutr* 1997;51:417
- Delzenne N, Roberfroid M. Physiological effects of non digestible oligosaccharides. *Lebensm Wiss U Technol* 1994;27:1
- Hidaka H, Eida T, Takizawa T, Tokunaga T, Tashiro Y. Effects of fructooligosaccharides on intestinal flora and human health. *Bifidobact Microflora* 1986;5:37
- Ziesenitz SC, Siebert G. In vitro assessment of nystose as a sugar substitute. *J Nutr* 1987;117:846
- Molis C, Florie B, Ouarne F, et al. Digestion, excretion, and energy value of fructooligosaccharides in healthy humans. *Am J Clin Nutr* 1996;64:324
- Knudsen KEB, Hessov I. Recovery of inulin from Jerusalem artichoke (*Helianthus tuberosus* L.) in the small intestine of man. *Br J Nutr* 1995;74:101
- Ellegard L, Andersson H, Bosaeus I. Inulin and oligofructose do not influence the absorption of cholesterol, and the excretion of cholesterol, Fe, Ca, Mg and bile acids but increases energy excretion in man. A blinded controlled cross-over study in ileostomy subjects. *Eur J Clin Nutr* 1997;51:1
- Wise EC, Heyl FW. Failure of a diabetic to utilize inulin. *J Am Pharm Soc Bull* 1931;20:26
- Alles MS, Hautvast JGA, Nagengast FM, et al. Fate of fructo-oligosaccharides in the human intestine. *Br J Nutr* 1996;76:211
- Gibson GR, Roberfroid MB. Dietary modulation of the human colonic microbiota—introducing the concept of prebiotics. *J Nutr* 1995;125:1401
- McBain A, Macfarlane G. Investigations of bifidobacterial ecology and oligosaccharide metabolism in a three-stage compound continuous culture system. *Scand J Gastroenterol* 1997;222(suppl):32
- Buddington RK, Williams CH, Chen S, Witherly SA. Dietary supplement of neosugar alters and fecal flora and decreases activities of some reductive enzymes in human subjects. *Am J Clin Nutr* 1996;63:709
- Gibson GR, Beatty ER, Wang X, Cummings JH. Selective stimulation of bifidobacteria in the human colon by oligofructose and inulin. *Gastroenterology* 1995;108:975
- Bouhnik Y, Flourie B, Riottot M, et al. Effects of fructo-oligosaccharides ingestion on fecal bifidobacteria and selected metabolic indexes of colon carcinogenesis in healthy humans. *Nutr Cancer* 1996;26:21
- Kleessen B, Sykura B, Zunft HJ, Blaut M. Effects of inulin and lactose on fecal microflora, microbial activity, and bowel habit in elderly constipated persons. *Am J Clin Nutr* 1997;65:1397
- Roberfroid MB, Van Loo JAE, Gibson GR. The bifidogenic nature of chicory inulin and its hydrolysis products. *J Nutr* 1998;128:11
- Gibson GR. Dietary modulation of the human gut microflora using the prebiotic inulin and oligofructose. *J Nutr* 1999;129(suppl 7S):1438S
- Rao AV. Dose-response effects of inulin and oligofructose on intestinal bifidogenesis effect. *J Nutr* 1999;129:1442S
- Hunter JO, Truffnell Q, Lee AJ. Controlled trial of oligofructose on the management of irritable bowel syndrome. *J Nutr* 1999;129(suppl 7S):1451S
- Catala I, Butel MJ, Bensada M, et al. Oligofructose contributes to the protective role of bifidobacteria in experimental necrotizing enterocolitis in quails. *J Med Microbiol* 1999;48:89
- Campbell JM, Fahey GJ, Wolf BW. Selected indigestible oligosaccharides affect large bowel mass, cecal and fecal short-chain fatty acids, pH and microflora in rats. *J Nutr* 1997;127:130
- Levrat MA, Remy C, Demigné C. High propionic acid fermentations and mineral accumulation in the cecum of rats adapted to different levels of inulin. *J Nutr* 1990;121:1730
- Remy C, Behr SR, Levrat M, Demigné C. Fiber fermentability in the rat cecum and its physiological consequences. *Nutr Res* 1992;12:1235
- Roland N, Nugon-Baudon L, Andrieux C, Szlyt O. Comparative study of the fermentative characteristics of inulin and different types of fiber in rats inoculated with a human whole fecal flora. *Br J Nutr* 1995;74:239
- Younes H, Demigné C, Remy C. Acidic fermentation in the caecum increases absorption of calcium and magnesium in the large intestine of the rat. *Br J Nutr* 1996;75:301
- Roberfroid MB, Delzenne N. Dietary fructans. *Annu Rev Nutr* 1998;18:117
- Van Loo J, Franck A, Cummings JH, et al. Consensus document on the functional food properties of non digestible oligosaccharides (DGXII ENDO Project). *Br J Nutr* 1999;81:121
- Delzenne N, Aertssens J, Verplaete H, Roccaro M, Roberfroid M. Effect of fermentable fructo-oligosaccharides on mineral, nitrogen and energy digestive balance in the rats. *Life Sci* 1995;57:1579
- Ohta A, Ohtsuki M, Baba S, et al. Calcium and magnesium absorption from the colon and rectum are increased in rats fed fructooligosaccharides. *J Nutr* 1995;125:2417
- Trautwein EA, Rieckhoff D, Erbersdobler HF. Dietary inulin lowers plasma cholesterol and triacylglycerol and alters biliary bile acid profile in hamsters. *J Nutr* 1998;128:1937
- Van den Heuvel EGHM, Muys T, van Dokkum W, Schaafsma G. Oligofructose stimulates calcium absorption in adolescents. *Am J Clin Nutr* 1999;69:544
- Coudray C, Bellange J, Castiglia-Delahauc C, et al. Effect of soluble and partly soluble dietary fibres supplementation on absorption and balance of calcium, magnesium, iron and zinc in healthy young men. *Eur J Clin Nutr* 1997;51:375
- Reddy BS, Hamid R, Rao CV. Effect of dietary oligofructose and inulin on colonic preneoplastic aberrant crypt foci inhibition. *Carcinogenesis* 1997;18:1371
- Rowland IR, Rumney CJ, Coutts JT, Lievens LC. Effect of *Bifidobacterium longum* and inulin on gut bacterial metabolism and carcinogen-induced aberrant crypt foci in rats. *Carcinogenesis* 1998;19:281
- Roberfroid MB. Caloric value of inulin and oligofructose. *J Nutr* 1999;129(suppl 7S):1436S
- Carabin I, Flamm G. Evaluation of safety of inulin, and oligofructose as dietary fiber. *Regul Tox Pharm* 2000(in press)
- Clevenger MA, Turnbull D, Inoue H, et al. Toxicological evaluation of neosugar: genotoxicity, carcinogenicity, and chronic toxicity. *J Am Coll Toxicol* 1988;7:643
- Briet F, Achour L, Flourie B, et al. Symptomatic response to varying levels of fructo-oligosaccharides consumed occasionally or regularly. *Eur J Clin Nutr* 1995;49:501
- Garleb KA, Snook JT, Marcon MJ, Wolf BW, Johnson CD. Effect of fructooligosaccharide containing enteral formulas on subjective tolerance factors, serum chemistry profiles, and faecal bifidobacteria in healthy adult male subjects. *Microbiol Ecol Health Dis* 1996;9:279
- Den Hond EM, Geypens BJ, Ghoos YF. Effect of long chain chicory inulin on bowel habit and transit time in constipated persons. *Nutr Res* 2000(in press)
- Trowell HC, Southgate DAT, Wolever TMS, et al. Dietary fiber redefined. *Lancet* 1976;i:967

Fructooligosaccharides from chicory (FOSC) are functional prebiotic foods recognized to exert several well-being effects in human health and animal production, as decreasing blood lipids, modulating the gut immune system, enhancing mineral bioavailability, and inhibiting microbial growth, among others. Mechanisms of actions directly on cell metabolism and structure are however little known. View at: Google Scholar. M. B. Roberfroid, "Chicory fructooligosaccharides and the gastrointestinal tract," *Nutrition*, vol. 16, no. 7-8, pp. 677-679, 2000. View at: Publisher Site | Google Scholar. E. Biedrzycka and M. Bielecka, "Prebiotic effectiveness of fructans of different degrees of polymerization," *Trends in Food Science & Technology*, vol. 15, no. 3-4, pp. 170-175, 2004. PDF | Chicory (*Cichorium intybus*) is a perennial herbal plant of the dandelion family Asteraceae, usually with bright blue flowers, rarely pink or | Find, read and cite all the research you need on ResearchGate. Fructooligosaccharides, which can be used to manipulate the composition of microbiota in the gastrointestinal tract and improve its integrity. Inulin is one of the best sources of fructooligosaccharides. Fructooligosaccharides are basically obtained from plant sources and comprise 3-10 fructose units with the terminal fructose linked to a glucose residue by 1-2 glycosidic bonds, that is, they have a nonreducing sucrose end. FOS and inulin are classified based on their DP, which is based on the number of monosaccharide units. FOS have a DP <10, while inulin has a DP ranging between 2 and 60. Most subjects do not tolerate FOS levels greater than 30 g/day due to adverse gastrointestinal symptoms [84]. Some studies have not observed any changes in blood lipids and lipoproteins with FOS intakes ranging between 9 and 20 g/day [85, 86], while others report significant reductions in total and LDL cholesterol or TGs [87, 88].