



Review Paper

Pinta–A Treponemal Infection to the Skin

Bincy Varghese* and Jayakrishnan S.S.

Department of Pharmacy Practice, College of Pharmaceutical Sciences, Govt. Medical College, Thiruvananthapuram, Kerala, 695011, INDIA
binzvarghese90@gmail.com

Available online at: www.isca.in, www.isca.me

Received 18th June 2015, revised 5th January 2016, accepted 20th January 2016

Abstract

Pinta is the oldest and mild form of the treponemal infections caused by Treponema carateum. The word pinta means painted and it was originated from Spanish. Lesions, red to bluish-black colored spots, splotches and discoloration of the skin are the major manifestations of this skin related bacterial infection. Reports suggest that pinta has virtually disappear from countries of Latin America and is now restricted to the Philippines and some areas of the Pacific. The following aspects of pinta such as its origin, prevalence and mode of transmission along with the clinical conditions, diagnostic methods, treatments and preventive measures are briefed in this review.

Keywords: Bacterial, Carateum, Infection, Pinta, Skin Lesions, Splotches, Spots, Treponema.

Introduction

Synonyms: Azul, Carate, Empeines, Lota, Mal de Pinto, Tina, Purupuru.

Pinta, a rare infectious tropical disease affecting the skin is caused by the bacterium *Treponema pallidum carateum* which is morphologically and serologically indistinguishable from the organism that causes syphilis¹. It is a human skin disease endemic to Mexico, Central America and South America. This disease is classified as a treponematoses, because it is an infectious disease caused by treponemas (a genus of spiral-shaped bacteria (spirochetes)).

The infection is acquired in childhood or early adolescence among people living in unhygienic conditions preferably in males. Pinta progresses through three distinct stages, which are characterized by various skin lesions and discoloration. The exposed lower extremities on the legs and ankles (65-85% of cases), the buttocks, arms, hands and face are the most common areas affected with primary lesions. The lesions remain active and infectious for many years leading to extensive depigmentation and acral dyschromia. Disorders associated with dyspigmentation such as tinea versicolor, vitiligo, melasma, leprosy are often in the differential diagnosis of pinta. Other organ systems are not affected. Exposed areas of the skin such as the face and extremities are most often affected.

Epidemiology

Rural, poverty-stricken areas of northern South America, Mexico and the Caribbean have been reported with the primary incidence of pinta. About 1 million cases of pinta were reported in Central and South America in 1950. 20% seropositivity is

found in remote rural areas of Panama in 1980's. No cases are yet reported in the United States. The current prevalence of pinta is unknown, but only a few hundred cases have been reported per year¹. Data on prevalence of pinta are limited but it might remain endemic in remote regions of Mexico (states of Oaxaca, Guerrero, Michoacan and Chiapas) where it was common in 1980's. Among Indian tribes in the Amazon region of Brazil, Colombia and Peru² has also reported with this disease.

Pinta – At A Glance

Feature	Pinta
Organism	T. carateum
Modes of transmission	Skin-to-skin
Usual age of acquisition	Late childhood
Primary lesion	Non ulcerating papule with satellites, pruritic
Location	Extremities, face
Secondary lesion	Pintides, pigmented pruritic
Infectious relapses	None
Late complications	Non destructive, dyschromic macules.

Mode of Transmission

This disease is believed to be transmitted by repeated direct person to person skin contact. Treponemas are abundant in early lesions and persist through to the late dyschromic stage. The

bacteria enter the skin through a small cut, scratch or other skin damage. High atmospheric humidity intensify the lesions, increasing their oozing and infectiousness³.

Reservoir

Those aged 15-30 years with long-standing skin lesions comprise the main reservoir.

Presentation

The least severe of treponemal infections pinta confined to the skin, is transmitted by infection-to-skin contact and produces a raised papule which enlarges and becomes hyperkeratotic (scaly/flaky) after an incubation period of two to three week. The exposed surface of arms and legs produces the lesions followed by local lymph node enlargement. Further thickening and appearance of flat lesions (pintids) all over the body occurs after three to nine months which are generally resolved. In certain cases a minor affected proportion may get worsened into the development of an end stage disease, pronounced as widespread pigmentary change with a mixture of hyper pigmentation and depigmentation which can be disfiguring⁴.

Symptoms of Pinta

Skin gets only affected by pinta which is initiated by flat, itchy, reddened areas on the hands, feet, legs, arms, face or neck. Several months later the same areas on both sides of the body preferentially where bones are close to the skin appears with slate-blue patches. The patches later on lose their colour and thickening of the affected skin on the palms and soles occurs.

Diagnosis

Clinical diagnosis is usually done, but as in case of yaws and bejel serological tests for syphilis, such as rapid plasma reagin (RPR), VDRL (venereal disease research laboratory) test and TPHA (T. pallidum haemagglutination), will be positive and on dark field microscopy of samples taken from the early papules are seen with spirochetes.

Treatment

The disease can either be treated with drugs such as penicillin, tetracycline, azithromycin, chloramphenicol or can be prevented through contact tracing by public health officials. Endemic treponematoses including pinta, yaws and bejel can be effectively treated by a long acting single intramuscular injection of penicillin⁵. It is usually treated very effectively with a single intramuscular injection of Benzathine penicillin G (Bicillin) in the upper outer quadrant of the buttock. The dose is 600,000 units for children and contacts under 6 years of age. 1.2 million units for those aged 6-15 years and 2.4 million units for adults. Alternate antibiotics can be given to those who are allergic to penicillin. Erythromycin or tetracycline 500mg by mouth four times a daily for 15 days is recommended for those

allergic to penicillin. Children between the ages of 8-15 years should receive half the dose. Tetracycline should not be given to pregnant women and children under 12 years of age^{6,7}.

The treatment policies recommended by WHO are as follows: If the prevalence of clinically active infection in the community is over 10%, give of Benzathine penicillin G to the entire population. If the prevalence of clinically active infection cases are 5-10%, give Benzathine penicillin G to the patients, their contacts and to all children below the age of 15 years. If the prevalence of clinically active infection is under 15% treat all active cases as well as households and other obvious contacts with Benzathine penicillinG⁸.

Complications and Sequelae of Pinta

Chancre, VDRL positive, Hypo pigmentation, Gumma, Skin ulceration and Facial ulcers etc.

Prevention

In pinta an attack on social and economic conditions of life is as important as an attack on the biological cause. Recrudescence of the disease is apt to occur unless environmental improvement is promoted⁹. For example: Improvement of personnel and domestic hygiene. Adequate water supply. Liberal use of soaps. Better housing conditions. Improvement of quality of life. Promoting community hygiene. Conducting various health education programmes. People residing in the areas more prone to pinta and travelers to such areas should avoid skin to skin contact with the infected people. People residing in the affected area should use insect repellants, mosquito nets etc as insect bite aids in the spread of pinta. Should keep the houses clean and should not leave stagnant water as it will attract mosquitoes and other insects. Sustained surveillance, integrated into strengthened primary health care must be maintained to detect and treat new or missed cases including their contacts and treatment failures.

Conclusion

It can be concluded that there is a need to conduct various health education and awareness programs for the proper and complete eradication of the disease. The standards of living, personnel and environmental hygiene must be improved. Equitable access to pinta prevention and treatment should be addressed early in the eradication effort.

References

1. Pecher S.A. and Croce J. (1988), Immunology of tertiary pinta. *Med Cutan Ibero Lat Am*, 16, 111-114.
2. Hopkins D.R. and Florez D. (1977). Pinta, yaws and general syphilis in Colombia. *Int J Epidemiology*, 6, 349-355.
3. Hackett C.J. (1953). Extend and nature of the yaws

- problem in Africa. *Bull World Health Organisation*, 8, 129-182.
4. Torok E. (2009). *Oxford Handbook of Infectious Diseases and Microbiology* (first ed.). Oxford University Press. 388. ISBN 978 0 19 856925 1.
 5. Fine Steven (2014). *Treponematoses (Endemic Syphilis) Medication*. Retrieved 15 September 2014.
 6. *Manson's Tropical Diseases* (2009). Edited by Gordon C Cook and Almuddin I Zumla; 22nd edition; chapter 68 Endemic Treponematoses, page no: 1146-1147.
 7. *Harrison's Principles of Internal Medicine* (2015). Longo, Fauci, Hauser, Kasper, Jameson, Loscalzo: Vol I. chapter 154; Endemic Treponematoses: page no:1389-1390.
 8. World Health Organization (1982). *Treponemal infection, World Health Organ Tech Rep Ser*, 674, 16-20.
 9. *Park's Textbook of Preventive and Social Medicine* (2011); K Park. 21st edition; Endemic Treponematoses Pinta; Page no: 314.

Pinta is a skin infection caused by the bacterium *Treponema carateum*, a relative of the bacterium which causes syphilis. The word "pinta" comes from the Spanish and means "painted." Pinta is also known as "azula" (blue), and "mal de pinto" (pinto sickness). It is one of several infections caused by different *Treponema* bacteria, which are called "endemic" or "non-venereal" treponematoses. A treponemal infection characterized by bizarre pigmentary changes in the skin occurring in tropical America; it is effectively treated by penicillin. pinta fever a disease observed in northern Mexico, identical with Rocky Mountain spotted fever. Miller-Keane Encyclopedia and Dictionary of Medicine, Nursing, and Allied Health, Seventh Edition. Pinta is a skin disease caused by a bacterial infection. It is usually acquired during childhood and contracted through the skin to skin contact with an infected person. The disease is endemic to Mexico, Central America and South America. About One million of cases with Pinta were reported in Central and South America in the year 1950. The word "Pinta" is a Spanish word which means "painted". Pinta is classified under a treponemal disease because it is contagious and is caused by treponemes, a genus of spiral-shaped bacteria. Recent reports suggest the occurrence of disease in the Philippines and some areas of the pacific region. The bacterium enters the skin through a cut, scratch or a lesion and causes a red scaly bump called the primary lesion. Pinta (also known as azul, carate, empeines, lota, mal del pinto, and tina) is a human skin disease caused by infection with the spirochete, *Treponema carateum*, which is morphologically and serologically indistinguishable from the bacterium that causes syphilis. The disease is endemic to Mexico, Central America, and South America. Pinta, the least severe of treponemal infections being limited to the skin, is thought to be transmitted by skin-to-skin contact (similar to bejel and yaws), and after an... Pinta, the least severe of treponemal infections being limited to the skin, is thought to be transmitted by skin-to-skin contact (similar to bejel and yaws), and after an incubation period of two to three weeks, produces a raised papule, which enlarges and becomes hyperkeratotic (scaly/flaky). Lesions are usually present in the exposed surface of arms and legs. Local lymph nodes might be enlarged. Three to nine months later, further thickened and flat lesions (pintids) appear all over the body. These generally resolve, but a proportion of people with pinta will go on to develop late-stage disease, characterised by widespread pigmentary change with a mixture of hyperpigmentation and depigmentation which can be disfiguring.[2]. Cause. Pinta is caused by the bacterium *Treponema carateum*.