

## Carcinogenicity of shift-work, painting, and fire-fighting

Kurt Straif, Robert Baan, Yann Grosse, Béatrice Secretan, Fatiha El Ghissassi, Véronique Bouvard, Andrea Altieri, Lamia Benbrahim-Tallaa, Vincent Coglianò, on behalf of the WHO International Agency for Research on Cancer Monograph Working Group

In October, 2007, 24 scientists from ten countries met at the International Agency for Research on Cancer (IARC), Lyon, France, to assess the carcinogenicity of shift-work, painting, and fire-fighting. These assessments will be published as volume 98 of the IARC Monographs.<sup>1</sup>

About 15–20% of the working population in Europe and the USA is engaged in shift-work that involves nightwork, which is most prevalent (above 30%) in the health-care, industrial manufacturing, mining, transport, communication, leisure, and hospitality sectors. Among the many different patterns of shift-work, those including nightwork are the most disruptive for the circadian clock.

Six of eight epidemiological studies from various geographical regions, most notably two independent cohort studies of nurses engaged in shift-work at night,<sup>2,3</sup> have noted a modestly increased risk of breast cancer in long-term employees compared with those who are not engaged in shiftwork at night. These studies are limited by potential confounding and inconsistent definitions of shift-work, with several focused on a single profession. The incidence of breast cancer was also modestly increased in most cohorts of female flight attendants,<sup>4</sup> who also experience circadian disruption by frequently crossing time zones. Limitations of studies in these flight attendants include the potential for detection bias, proxy measures of exposure, and potential uncontrolled confounding by reproductive factors and cosmic radiation.

Several different rodent models have been used to test the effect of disruption of the circadian system

on tumour development. More than 20 studies investigated the effect of constant light, dim light at night, simulated chronic jet lag, or circadian timing of carcinogens, and most showed a major increase in tumour incidence. No clear effect was seen for light pulses at night or constant darkness. A similar number of studies investigated the effect of reduced nocturnal melatonin concentrations or removal of the pineal gland (where melatonin is produced) in tumour development and most showed increases in the incidence or growth of tumours.<sup>5,6</sup>

Exposure to light at night disturbs the circadian system with alterations of sleep-activity patterns, suppression of melatonin production, and deregulation of circadian genes involved in cancer-related pathways.<sup>7</sup> Inactivation of the circadian Period gene, *Per2*, promotes tumour development in mice,<sup>8</sup> and in human breast and endometrial tumours, the expression of PERIOD genes is inhibited.<sup>9</sup> In animals, melatonin suppression can lead to changes in the gonadotrophin axis.<sup>10</sup> In humans, sleep deprivation and the ensuing melatonin suppression lead to immunodeficiency.<sup>11,12</sup> For example, sleep deprivation suppresses natural killer-cell activity<sup>13</sup> and changes the T-helper 1/T-helper 2 cytokine balance, reducing cellular immune defence and surveillance.<sup>14</sup>

On the basis of “limited evidence in humans for the carcinogenicity of shift-work that involves nightwork”, and “sufficient evidence in experimental animals for the carcinogenicity of light during the daily dark period (biological night)”, the Working Group concluded that “shift-work that involves circadian

disruption is probably carcinogenic to humans” (Group 2A).<sup>15</sup>

Painters are potentially exposed to many chemicals used as pigments, extenders, binders, solvents, and additives. Painters can also be exposed to other workplace hazards, such as asbestos or crystalline silica.

Cohort and linkage studies of painters have shown consistent and significant increases in lung cancer compared with the general population. No information on tobacco smoking was available in the cohort studies; however, the increases are comparable to results from many case-control studies that controlled for smoking. A meta-analysis by the Working Group of all independent studies, including two recent large studies,<sup>16,17</sup> showed a significant excess risk of about 20% overall, and of 50% when the analysis was restricted to smoking-adjusted estimates from population-based case-control studies. Increased mortality from mesothelioma was consistently noted.<sup>18,19</sup>

Similarly, cohort and linkage studies showed consistent 20–25% increases in the occurrence of urinary bladder cancer in painters, and similar increases were noted in case-control studies that controlled for smoking.<sup>15,17,20</sup>

Although findings for lymphatic and haemopoietic cancers in painters were inconsistent, four of five case-control studies reported significant increases in childhood leukaemia associated with maternal exposure to paint. The risks tended to be greater when mothers were exposed before or during, rather than after, pregnancy,<sup>21,22</sup> and two studies showed some evidence of an increasing risk with increasing exposure.



### Upcoming meetings

February 5–12, 2008

Industrial and cosmetic dyes and related exposures

<http://monographs.iarc.fr/>

Overall, little information existed on specific work settings, and no particular agent could be identified from the available epidemiological studies as the cause of the increases in lung and urinary bladder cancers in painters.

Most cytogenetic studies on painters reported increased levels of genetic damage, such as chromosomal aberrations, micronucleus formation, sister chromatid exchange,<sup>23</sup> and DNA single-strand breaks. Several studies showed a dose-response association with duration of work. Stratified analyses by tobacco-smoking status generally showed similar results in smokers and non-smokers.

The Working Group concluded that there is “sufficient evidence in humans that occupational exposure as a painter causes cancers of the lung and urinary bladder”. Additionally, there is “limited evidence in humans”, mainly on the basis of studies of maternal exposure, that painting is associated with childhood leukaemia.

Overall, occupational exposure as a painter is “carcinogenic to humans” (Group 1).<sup>15</sup>

Firefighters are exposed to many toxic combustion products, including many known, probable or possible carcinogens. These intermittent exposures can be intense, and short-term exposures can be high for respirable particulate matter and for some carcinogens, notably benzene, benzo[a]pyrene, 1,3-butadiene, and formaldehyde.

Although increases in various cancers in fire-fighters compared with the general population have been noted in several studies, consistent patterns are difficult to discern due to the large variations of exposures. The Working Group updated a recent meta-analysis of cancer in firefighters.<sup>24</sup> For three types of cancer the relative risks were consistently increased and the average increase was significant: testicular cancer (all six studies

showed increased risks, average relative risk 1.5), prostate cancer (increased risks in 18 of 21 studies, average relative risk 1.3), and non-Hodgkin lymphoma (increased risks in five of six studies, average relative risk 1.2).

For intermittent, but intense, exposures to highly variable complex mixtures, conventional measures, such as years of employment or number of firefighting runs, can be poor surrogates for exposure. The available epidemiological studies are inherently limited by this issue.

Acute and chronic inflammatory respiratory effects noted in firefighters<sup>25</sup> would provide a plausible mechanism for respiratory carcinogenesis. Studies that assess genotoxic effects in firefighters are few and inconclusive.

On the basis of “limited evidence of carcinogenicity in humans”, the Working Group classified occupational exposure as a firefighter as “possibly carcinogenic to humans” (Group 2B).<sup>15</sup>

The IARC authors declared no conflicts of interest.

- 1 IARC. IARC monographs on the evaluation of carcinogenic risks to humans. Volume 98. Shift-work, painting and fire-fighting. Lyon: International Agency for Research on Cancer (in press).
- 2 Schernhammer ES, Laden F, Speizer FE, et al. Rotating night shifts and risk of breast cancer in women participating in the nurses' health study. *J Natl Cancer Inst* 2001; **93**: 1563–68.
- 3 Schernhammer ES, Kroenke CH, Laden F, Hankinson SE. Night work and risk of breast cancer. *Epidemiology* 2006; **17**: 108–11.
- 4 Megdal SP, Kroenke CH, Laden F, Pukkala E, Schernhammer ES. Night work and breast cancer risk: a systematic review and meta-analysis. *Eur J Cancer* 2005; **41**: 2023–32.
- 5 Anisimov VN, Baturin DA, Popovich IG, et al. Effect of exposure to light-at-night on life span and spontaneous carcinogenesis in female CBA mice. *Int J Cancer* 2004; **111**: 475–79.
- 6 Anisimov VN. The light-dark regimen and cancer development. *Neuro Endocrinol Lett* 2002; **23**: 28–36.
- 7 Stevens RG, Blask DE, Brainard GC, et al. Meeting report: the role of environmental lighting and circadian disruption in cancer and other diseases. *Environ Health Perspect* 2007; **115**: 1357–62.
- 8 Fu L, Pelicano H, Liu J, Huang P, Lee C. The circadian gene *Period2* plays an important role in tumor suppression and DNA damage response in vivo. *Cell* 2002; **111**: 41–50.

- 9 Chen ST, Choo KB, Hou MF, et al. Deregulated expression of the PER1, PER2 and PER3 genes in breast cancers. *Carcinogenesis* 2005; **26**: 1241–46.
- 10 Goldman BD. Mammalian photoperiodic system: formal properties and neuroendocrine mechanisms of photoperiodic time measurement. *J Biol Rhythms* 2001; **16**: 283–301.
- 11 Everson CA. Sustained sleep deprivation impairs host defense. *Am J Physiol* 1993; **265**: R1148–54.
- 12 Nelson RJ. Seasonal immune function and sickness responses. *Trends Immunol* 2004; **25**: 187–92.
- 13 Irwin M, McClintick J, Costlow C, Fortner M, White J, Gillin JC. Partial night sleep deprivation reduces natural killer and cellular immune responses in humans. *FASEB J* 1996; **10**: 643–53.
- 14 Dimitrov S, Lange T, Tiekens S, Fehm HL, Born J. Sleep associated regulation of T helper 1/T helper 2 cytokine balance in humans. *Brain Behav Immun* 2004; **18**: 341–48.
- 15 IARC. IARC monographs on the evaluation of carcinogenic risks to humans. Preamble. Lyon, France: International Agency for Research on Cancer, 2006.
- 16 Steenland K, Palu S. Cohort mortality study of 57,000 painters and other union members: a 15 year update. *Occup Environ Med* 1999; **56**: 315–21.
- 17 Bouchardy C, Schuler G, Minder C, et al. Cancer risk by occupation and socioeconomic group among men—a study by the Association of Swiss Cancer Registries. *Scand J Work Environ Health* 2002; **28**: 1–88.
- 18 Peto J, Hodgson JT, Matthews FE, Jones JR. Continuing increase in mesothelioma mortality in Britain. *Lancet* 1995; **345**: 535–39.
- 19 Brown LM, Moradi T, Gridley G, et al. Exposures in the painting trades and paint manufacturing industry and risk of cancer among men and women in Sweden. *J Occup Environ Med* 2002; **44**: 258–64.
- 20 Silverman DT, Levin LI, Hoover RN, Hartge P. Occupational risks of bladder cancer in the United States: I. White men. *J Natl Cancer Inst* 1989; **81**: 1472–80.
- 21 Shu XO, Stewart P, Wen WQ, et al. Parental occupational exposure to hydrocarbons and risk of acute lymphocytic leukemia in offspring. *Cancer Epidemiol Biomarkers Prev* 1999; **8**: 783–91.
- 22 Schüz J, Kaletsch U, Meinert R, Kaatsch P, Michaelis J. Risk of childhood leukemia and parental self-reported occupational exposure to chemicals, dusts, and fumes: results from pooled analyses of German population-based case-control studies. *Cancer Epidemiol Biomarkers Prev* 2000; **9**: 835–38.
- 23 Testa A, Festa F, Ranaldi R, et al. A multi-biomarker analysis of DNA damage in automobile painters. *Environ Mol Mutagen* 2005; **46**: 182–88.
- 24 LeMasters GK, Genaidy AM, Succop P, et al. Cancer risk among firefighters: a review and meta-analysis of 32 studies. *J Occup Environ Med* 2006; **48**: 1189–202.
- 25 Burgess JL, Brodtkin CA, Daniell WE, et al. Longitudinal decline in measured firefighter single-breath diffusing capacity of carbon monoxide values. A respiratory surveillance dilemma. *Am J Respir Crit Care Med* 1999; **159**: 119–24.

**Monograph Working Group Members**

A Blair—Chair (USA); P Demers (Canada); J Hansen (Denmark); E Pukkala (Finland); F Lévi (not present for evaluations; France); T Brock (Germany); G Costa, F Merletti (Italy); R Vermeulen (Netherlands); M Bråtveit (Norway); D Blask, J Burgess, S Davis, E Haus, P Landrigan (unable to attend), G LeMasters, C Portier, E Schernhammer, K Steenland, R Stevens, T Zheng, Y Zhu (USA)

**Conflicts of interest**  
None

**Invited Specialists**  
J Arendt, J Cherie (UK); C Austin (USA)

**Conflicts of interest**  
JA has received unrestricted research funding from Philips Lighting (Eindhoven, Netherlands), which was previously a joint patent holder for the use of short-wavelength light for adjusting circadian rhythms and maintaining alertness. JA is the Director of Stockgrand Ltd (Surrey, UK), which sells melatonin-measurement and 6-sulphatoxymelatonin-measurement technology. Finally, JA is a consultant at Alliance Pharmaceuticals UK, which develops melatonin products. JC undertakes consulting work for paint and pigment manufacturers, funded by European chemical industry trade associations and large industrial companies.

**Representative**  
A Huici-Montagud (European Commission)

**Observer**  
K Mundt (International Paint and Printing Ink Council)

Shift work with circadian disruption has been considered as a carcinogenic risk factor for skin cancer. The few prior studies that investigated the association between shift work and skin cancer have inconclusive results. Our main objective was to evaluate the associations between shift work and the risks of different types of skin cancer. We systematically searched PubMed, Web of Science, Cochrane Library, EMBASE and Science Direct until October 2018 for studies that included a relationship between shift work and skin cancer. Our search yielded 193 articles and 9 studies met the criteria for our review.Â Straif, K. et al. Carcinogenicity of shift-work, painting, and fire-fighting. *Lancet Oncol.* 8, 1065â€“1066 (2007). Shift work and chronobiology have a longstanding tradition of successful research behind them, particularly in Germany, where Rutenfranz and Aschoff were among the prominent figures in this area. Nonetheless, the question whether shift work contributes to the development of cancer by way of circadian disruption or chronodisruption is very new, and relatively few studies have been carried out with a view toward practical preventive measures.Â Carcinogenicity of shift-work, painting, and fire-fighting. *Lancet Oncol* 2007; 8: 1065â€“6. MEDLINE. Asthma and other adverse health effects of isocyanates are well-documented and exposure surveillance is crucial to disease prevention. Hexamethylene diisocyanate (HDI)-specific serum immunoglobulin G (IgG) was evaluated as an exposure biomarker among workers at a US Air Force Air Logistics Center, which includes a large aircraft maintenance facility.Â *Int J Occup Med Environ Health* 2010 ;23(4):331-8. Department of Public Health and Preventive Medicine, Faculty of Medicine, The University of Mansoura, Mansoura, Egypt. Objectives: This study examined the association between male infertility and certain occupational exposures. Material And Methods: A case-control study was carried out from January 2008 to February 2009; on 255 infertile men and 267 fertile men controls. Painting, Firefighting, and Shiftwork. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 98. IARC.Â This volume of the IARC Monographs provides evaluations of the carcinogenicity of shiftwork, painting and firefighting. Shiftwork is estimated to involve about 15-20% of the total working population. It is most prevalent among workers in the health care, transportation, communication, leisure and hospitality sectors. Shiftwork involving work at night is the most disruptive for the circadian clock. Painters are potentially exposed to the chemicals found in paint products during their application and removal, and may also be exposed to other workplace hazards, such as asbestos or crystalline silica dust. Straif, K., Baan, R., Grosse, Y., Secretan, B., Ghissassi, F., Bouvard, V., Altieri, A., Benbrahim-Tallal, L. and Coglianò, V. (2007) Carcinogenicity of Shift-Work, Painting, and Fire-Fighting. *Lancet Oncology*, 8, 1065-1066. [http://dx.doi.org/10.1016/S1470-2045\(07\)70373-X](http://dx.doi.org/10.1016/S1470-2045(07)70373-X). has been cited by the following articleÂ The odds ratio of working night shift was found to be 1.24 (OR = 1.24, p = 0.615) higher in breast cancer research participants compared to research participants diagnosed with other types of cancerâ€”odds ratio of 0.80 (p = 0.610). For rotational work, the OR was 1.445, indicating a higher risk than for shift work. It is recommended that the relationship between working night shift and breast cancer risk be explored further through cross-sectional and cohort studies.